

*Degradation of N-Nitrosodimethylamine  
by UV-Based Advanced Oxidation  
Processes for Potable Reuse: a Short Review*

**Takahiro Fujioka, Shunya Masaki,  
Hitoshi Kodamatani & Keisuke Ikehata**

**Current Pollution Reports**


e-ISSN 2198-6592

Curr Pollution Rep  
DOI 10.1007/s40726-017-0052-x



**Your article is protected by copyright and all rights are held exclusively by Springer International Publishing AG. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at [link.springer.com](http://link.springer.com)".**

# Degradation of *N*-Nitrosodimethylamine by UV-Based Advanced Oxidation Processes for Potable Reuse: a Short Review

Takahiro Fujioka<sup>1</sup>  · Shunya Masaki<sup>1</sup> · Hitoshi Kodamatani<sup>2</sup> · Keisuke Ikehata<sup>3</sup>

© Springer International Publishing AG 2017

**Abstract** The ultraviolet (UV)-based advanced oxidation process (AOP) is a powerful technology commonly utilised in recent potable water reuse (PR) schemes. The AOP involves the generation of highly reactive free radicals (e.g. hydroxyl, HO<sup>•</sup>) and is primarily applied for the removal of two target trace organic chemicals—*N*-nitrosodimethylamine (NDMA) and 1,4-dioxane—in the PR schemes. Both of these organics are not well removed by the reverse osmosis (RO) process. NDMA is a probable carcinogen and is often present in reclaimed water at concentrations higher than the guidelines established for PR. This review aimed to provide an understanding of the current UV-based advanced oxidation technologies for NDMA removal in PR, their limitations and the future of advanced technologies for their removal. NDMA is readily photolysed by direct UV irradiation, while an AOP such as UV/H<sub>2</sub>O<sub>2</sub> process is necessary for the destruction of 1,4-dioxane. Unfortunately, the generation of hydroxyl radicals through UV photolysis of H<sub>2</sub>O<sub>2</sub> is largely inefficient with conversion on the order of 20% under normal plant operations and the addition of H<sub>2</sub>O<sub>2</sub> (e.g. 3 mg/L) provides only a negligible improvement in NDMA destruction. However, AOP can

also be achieved without continuous chemical addition through the application of UV irradiation to heterogeneous photocatalysts (e.g. TiO<sub>2</sub>). The UV/TiO<sub>2</sub> process generates hydroxyl radicals and singlet oxygen molecules, both of which degrade NDMA into by-products (e.g. methylamine or dimethylamine). Recent studies revealed that modification of the surface morphology of TiO<sub>2</sub> can not only enhance NDMA destruction but also alter the composition of the degradation by-products.

**Keywords** *N*-nitrosodimethylamine (NDMA) · Hydroxyl radicals · Photocatalytic degradation · Potable reuse · Titanium oxide · 1,4-Dioxane

## Introduction

The occurrence and fate of *N*-nitrosodimethylamine (NDMA; C<sub>2</sub>H<sub>6</sub>N<sub>2</sub>O, molecular weight 74 g/mol) in potable water reuse (PR) have attracted significant attention in recent years [1, 2]. NDMA is an *N*-nitrosamine that has been classified as probable carcinogens by the US EPA [3]. Although reverse osmosis (RO) membrane treatment in water reclamation systems is a key barrier for the elimination of trace organic chemicals (TrOCs) in wastewater [4], NDMA readily permeates through RO due to its small size and non-ionic nature under treatment conditions. NDMA rejection by RO membranes has been reported at 10–70% [5–7]. The other challenging carcinogenic TrOC, 1,4-dioxane (C<sub>4</sub>H<sub>8</sub>O<sub>2</sub>, molecular weight 88 g/mol), has an RO rejection similar to NDMA [8]. To comply with the regulatory guidelines and meet the PR notification levels of NDMA (10 ng/L) [9, 10] and 1,4-dioxane (1 µg/L) [11], most PR facilities in the USA and Australia are equipped with a post treatment–ultraviolet (UV)/hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) advanced oxidation process (AOP) [7, 12].

---

This article is part of the Topical Collection on *Water Pollution*

---

✉ Takahiro Fujioka  
tfujioka@nagasaki-u.ac.jp

- <sup>1</sup> Water and Environmental Engineering, Graduate School of Engineering, Nagasaki University, 1-14 Bunkyo-machi, Nagasaki 852-8521, Japan
- <sup>2</sup> Division of Earth and Environmental Science, Graduate School of Science and Engineering, Kagoshima University, 1-21-35 Korimoto, Kagoshima 890-0065, Japan
- <sup>3</sup> Pacific Advanced Civil Engineering, Inc., 17520 Newhope St., Suite 200, Fountain Valley, CA 92708, USA

The UV/H<sub>2</sub>O<sub>2</sub> AOP involves the use of hydroxyl radicals that form through UV photolysis of H<sub>2</sub>O<sub>2</sub> added to the reactor feedwater. Although NDMA is a photodegradable compound that can be readily destroyed by direct UV irradiation, 1,4-dioxane requires some form of radicals for its destruction. Inherent disadvantages of the UV/H<sub>2</sub>O<sub>2</sub> AOP include the requirement of a continuous dose of chemical oxidant (e.g. H<sub>2</sub>O<sub>2</sub>). Moreover, only about 20% of H<sub>2</sub>O<sub>2</sub> is consumed through the AOP [13, 14], meaning that the system requires a quenching step of residuals H<sub>2</sub>O<sub>2</sub> using sodium thiosulphate (e.g. 2.5–3 mg dose to 1 mg-H<sub>2</sub>O<sub>2</sub>/L) [13], sodium hypochlorite [15] or granular activated carbon [16] in the following step. The occurrence of 1,4-dioxane in treated wastewater is more site specific because it is an industrial solvent and is often detected at concentrations lower than the regulated values [17]. As a result, recent research focus has been placed on NDMA removal by AOPs.

Other AOPs that have been commercialised or extensively studied include UV/ozone [18] and ozone/H<sub>2</sub>O<sub>2</sub> [19, 20]. UV/ozone AOP is based on the formation of hydroxyl radicals through UV photolysis of ozone at wavelengths of 200–280 nm. Ozone/H<sub>2</sub>O<sub>2</sub> AOP generates hydroxyl radicals as a reaction product of the two oxidants (i.e. ozone and H<sub>2</sub>O<sub>2</sub>). Other emerging technologies include sulphate radical-based AOP [21, 22] and chlorine radical-based AOP [23, 24]. However, all of the AOPs described earlier require continuous dosing of some chemical. AOPs that do not require any chemical dosing for NDMA removal include UV irradiation of heterogeneous photocatalysts (e.g. titanium oxide, TiO<sub>2</sub>) [25, 26]. In UV/TiO<sub>2</sub> AOP, hydroxyl radicals are formed at the surface of photocatalysts (i.e. TiO<sub>2</sub>). Far less studies have been conducted for NDMA removal using UV/TiO<sub>2</sub> AOP. No large-scale UV/TiO<sub>2</sub> AOP system has been employed in any PR schemes to date.

In addition to the fate of NDMA in PR, the fate of NDMA precursors including dimethylamine (DMA) and tertiary amines has been extensively investigated [27, 28–30]. A considerable increase in NDMA can occur from NDMA precursors and the presence of residual chloramine [31–33], both of which often remain in the source water at the end of the treatment process. Hydroxyl radicals formed by UV/TiO<sub>2</sub> AOP can transform NDMA into precursors that could reform NDMA through the reaction with chloramine [27, 34]. The fate of NDMA precursors through UV/TiO<sub>2</sub> AOP is not very well understood. Therefore, a better understanding of the removal of NDMA and its precursors is a key factor to consider if UV/TiO<sub>2</sub> is to become a viable AOP.

This short review paper aimed to provide further understanding of the removal of NDMA and NDMA precursors through UV-based AOP with a particular focus on photocatalytic UV/TiO<sub>2</sub> AOP. The specific objectives are to clarify the effectiveness of UV-based AOP toward NDMA removal and to identify the degradation by-products. Thereafter, an

assessment of the feasibility of utilising UV/TiO<sub>2</sub> AOP in PR applications will be discussed.

## Occurrence and Fate of NDMA During Water Reclamation Processes

Municipal wastewater originates from a variety of domestic and industrial sources which contain a diverse range of constituents (e.g. dissolved and suspended solids, pathogens, organic and inorganic compounds). TrOCs are of great concern in PR, because they are difficult to remove by conventional water treatment technologies. These TrOCs include pharmaceuticals, personal care products, steroid hormones, pesticides, disinfection by-products and industrial chemicals. To ensure that safe and high-quality reclaimed wastewater is produced for the augmentation of drinking water supply, PR schemes are typically composed of a series of advanced water treatment processes that can remove a variety of constituents. These treatment processes include microfiltration (MF) or ultrafiltration (UF), RO and AOP [35–37]. MF/UF is employed as a pretreatment for RO to remove bacteria and suspended solids and minimise membrane fouling. RO treatment is a critical barrier against almost all constituents in water that include inorganics (ions and heavy metals) and most TrOCs. However, two specific TrOCs, NDMA (disinfection by-product) and 1,4-dioxane (solvent), readily permeate through RO membranes [38]. Both are uncharged and probably carcinogens [3, 39]. Among them, NDMA is often identified above the regulatory notification level (e.g. 10 ng/L in CA, USA) in the RO permeate at advanced water reclamation facilities [7]. To comply with the regulatory limits, an AOP (UV/H<sub>2</sub>O<sub>2</sub>) is typically applied following the RO treatment and therefore can successfully reduce NDMA concentrations to below its detection limit (i.e. 1–2 ng/L) [37, 40].



There are many sources of NDMA in treated wastewater. NDMA is typically present in primary wastewater effluent at the concentrations below 100 ng/L [41, 42]. In addition, NDMA forms through chloramination for mitigation of RO biofouling [5, 43], as well as during ozonation [44, 45]. DMA is widely reported as an NDMA precursor [31, 41], while others (e.g. amine-containing coagulation polymers, pharmaceuticals and personal care products, such as ranitidine—a component of antacids like Zantac®) have also been recognised to be its major sources [28, 43]. It should be noted that the molar conversion rate of DMA to NDMA is relatively low (<3% [46]) as compared with those of several tertiary amines that have DMA functional groups (e.g. ranitidine, 90% [28]). Most of the NDMA precursors are well removed by RO due to their large size, while some NDMA precursors can permeate through RO [47]. In addition, NDMA precursors can also be formed across the AOP through the degradation of NDMA and presence of residual chloramine in the effluent stream [27, 34].

### Degradation of NDMA by Direct UV

NDMA is readily degraded by direct photolysis at wavelengths <260 nm. It has a maximum absorbance at a wavelength of 227 nm, which originates from a  $\pi \rightarrow \pi^*$  transition, and absorbs UV light strongly at 254 nm ( $\epsilon = 1974 \text{ M}^{-1} \text{ cm}^{-1}$ ), which commercial low-pressure mercury-vapour UV lamps emit [48, 49]. NDMA also has a weak UV absorption at 332 nm due to an  $n \rightarrow \pi^*$  transition ( $\epsilon = 109 \text{ M}^{-1} \text{ cm}^{-1}$ ) [50]. Quantum yield of NDMA photolysis in a buffered solution with neutral pH (e.g. 7–8) and sufficient oxygen concentrations is >0.3 [48, 51]. A 1-log removal of NDMA can be achieved with  $1000 \text{ mJ/cm}^2$  UV dose [49].


A previous study conducted by Lee et al. [48] has suggested that UV degradation of NDMA in aqueous solution at pH <8.5 fundamentally undergoes three pathways: (1) homolytic cleavage of N–NO bonds, (2) heterolytic cleavage of N–NO bonds and (3) photooxidation. All the pathways start with the protonation of NDMA into an excited NDMA ( $\text{NDMA}^*-\text{H}^+$ ) following UV irradiation. In the first pathway, the excited NDMA decays to two species—aminium radical ( $(\text{CH}_3)_2\text{NH}^+$ ) and nitric oxide ( $\cdot\text{N O}$ ) (Eq. 1). They decay further to hyponitrous acid (HNO) and protonated *N*-methylidenemethylamine ( $\text{CH}_2 = \text{N}^+\text{HCH}_3$ ) [52]. The protonated *N*-methylidenemethylamine undergoes hydrolysis, producing methylamine ( $\text{CH}_3\text{NH}_3^+$ , MA) and formaldehyde (HCOH) (Eq. 2). In a neutral pH (e.g. pH >5.5) environment, the reaction between aminium radical ( $(\text{CH}_3)_2\text{NH}^+$ ) and  $\cdot\text{N O}$  leads to the formation of amidoxime ( $\text{CH}_3\text{NHC} = \text{NOH}$ ) (Eq. 3). The reaction between  $\text{CH}_3\text{NHC} = \text{NOH}$  and nitrite ion ( $\text{NO}_2^-$ ) forms *N*-methylformamide ( $\text{CH}_3\text{NCOH}$ ) and  $\text{N}_2\text{O}$ .

#### Pathway 1



In the second pathway, the excited NDMA reacts with a water molecule or  $\text{NO}_2^-$  that leads to heterolytic cleavage of N–NO bonds and the formation of DMA ( $(\text{CH}_3)_2\text{NH}_2^+$ ) and nitrite (Eq. 4) [51].


#### Pathway 2



In the third pathway, a one-electron transfer occurs from the excited NDMA and reacts with oxygen to yield protonated *N*-methylidenemethylamine ( $\text{CH}_2 = \text{N}^+\text{HCH}_3$ ),  $\cdot\text{N O}$  and

superoxide radical ( $\text{O}_2^{\cdot-}$ ) (Eq. 5). The protonated *N*-methylidenemethylamine ( $\text{CH}_2 = \text{N}^+\text{HCH}_3$ ) decays to MA and HCOH by hydrolysis (Eq. 2) [48]. The reaction between  $\cdot\text{N O}$  and  $\text{O}_2^{\cdot-}$  leads to the formation of peroxyxynitrite ( $\text{ONOO}^-$ ), which transforms into nitrate ion ( $\text{NO}_3^-$ ) (Eq. 6).

#### Pathway 3



DMA and MA are breakdown products associated with the degradation of NDMA by UV irradiation. Prolonged exposure to UV photolysis does not lead to a reduction in the concentration of DMA and MA [50, 53]. The major pathway of photolytic degradation is determined by the solution pH. At low pH (e.g. 3–4), pathway 2 is dominant with DMA and  $\text{NO}_2^-$  as major degradation products. DMA formation decreases with increasing pH, while MA formation increases. At water treatment pH 7–8, the pathways 1–3 concurrently occur leading to similar formation levels of DMA and MA. Overall, UV photolysis at higher pH can minimise the formation of DMA, which is advantageous because DMA is a major NDMA precursor. Nevertheless, NDMA photolysis at pH higher than approximately 8 could be negatively impacted by very low quantum yield [48, 51] due to the light scavenging effects of carbonate ions ( $\text{CO}_3^{2-}$ ), bicarbonate ions ( $\text{HCO}_3^-$ ) and dissolved organic matter (DOM) [53–55].

Selection of the UV lamp is an important factor for the destruction of NDMA [56]. Low-pressure (LP) mercury lamp emits monochromatic light at 254 nm, while the medium-pressure (MP) mercury lamp emits a broader spectrum of UV–visible light at wavelengths between 200 and 500 nm. Although the wide range of wavelengths of MP lamps covers the absorbance of many substances in the source water, it could lead to a lower energy efficiency for compounds with narrow absorbance spectrum. Sharpless et al. [49] reported that the photonic efficiencies (UV fluence-based rate constants,  $\text{cm}^2/\text{mJ}$  at wavelengths 200–300 nm) of LP and MP UV lamps were similar, with the MP lamp having an average light emission of approximately 256 nm between 200 and 300 nm. UV lamps with light emission ranges lower than LP and MP UV lamps (e.g. excimer UV lamps [57]) could be more suitable for NDMA removal due to the 227 nm absorbance maximum of NDMA. Sakai et al. [58] used a KrCl excimer lamp that emits monochromatic light at 222 nm and reported reaction rate constants in pure water at 11 and  $2.6 \text{ cm}^2/\text{J}$  for the KrCl excimer lamp and a LP lamp, respectively. The high efficiency of the KrCl excimer lamp was attributed to the high absorptivity and high quantum yield of NDMA at 222 nm [58]. KrCl excimer lamps have a significantly shorter lifetime (<4000 h) than LP and MP UV lamps (e.g. >10,000 h) [59].


## Degradation of NDMA by UV/H<sub>2</sub>O<sub>2</sub> AOP

AOP is a water treatment process that utilises strong oxidising agents (e.g. hydroxyl radical, HO<sup>•</sup>) to oxidise and degrade contaminants of public health concern [24<sup>•</sup>]. UV/H<sub>2</sub>O<sub>2</sub> AOP generates hydroxyl radicals through irradiation of H<sub>2</sub>O<sub>2</sub> at wavelengths <270 nm with a quantum yield of 1.11 [60]. UV irradiation photolyses H<sub>2</sub>O<sub>2</sub> ( $\epsilon_{245\text{ nm}} = 19.6\text{ M}^{-1}\text{ cm}^{-1}$ ) and generates hydroxyl radicals as follows:



Molar absorption coefficient of H<sub>2</sub>O<sub>2</sub> increases with decreasing UV wavelength. Because the molar absorption coefficient of H<sub>2</sub>O<sub>2</sub> is generally low at the wavelengths of typical UV lamp (e.g. 254 nm for LP mercury lamp), the production rate of hydroxyl radicals also remains low [13]. Accordingly, a high concentration of H<sub>2</sub>O<sub>2</sub> (3–5 mg/L) is required for H<sub>2</sub>O<sub>2</sub> to absorb more UV light [61].

Hydroxyl radical reactions include four basic pathways: (a) abstraction of a hydrogen atom usually for aliphatic hydrocarbon groups, (b) radical addition for unsaturated or aromatic hydrocarbon groups, (c) electron transfer for inorganics and (d) radical–radical reaction [61]. The degradation of NDMA by UV/H<sub>2</sub>O<sub>2</sub> AOP occurs by hydrogen atom abstraction from the methyl groups in NDMA as follows [49, 62]:



The hydroxyl radical rate constant for NDMA is  $4.3 \times 10^8\text{ M}^{-1}\text{ s}^{-1}$  [62], and 1-log removal of NDMA by UV/H<sub>2</sub>O<sub>2</sub> AOP can typically be achieved with a UV dose of  $\geq 1000\text{ mJ/cm}^2$  and 3 mg/L of H<sub>2</sub>O<sub>2</sub>. To the best of our knowledge, further degradation pathways of NDMA by UV/H<sub>2</sub>O<sub>2</sub> AOP have not well been established, while Lee et al. [19] have described the degradation mechanisms by hydroxyl radicals formed through ozone/H<sub>2</sub>O<sub>2</sub> AOP. In pathway 1, the carbon-centred NDMA radical self-decomposes to protonated *N*-methylidenemethylamine ( $\text{CH}_2 = \text{N}^+\text{HCH}_3$ ) (Eq. 9). The pathway 1 is a major pathway of NDMA degradation by hydroxyl radicals.




Pathway 1



In pathway 2, the carbon-centred NDMA radical reacts with dissolved oxygen to form a peroxy radical ( $\cdot\text{O}_2\text{CH}_2(\text{CH}_3)\text{N}-\text{N}=\text{O}$ ) (Eq. 10), which decomposes to protonated *N*-methylidenemethylamine ( $\text{CH}_2 = \text{N}^+\text{HCH}_3$ ) (Eq. 11). Both pathways 1 and 2 generate MA as the primary product from a reaction between NDMA and hydroxyl radicals (Eq. 12). The formed MA can be further degraded by hydroxyl radicals (rate constant  $k_{\text{HO}^\bullet} = 1.8\text{--}5.7 \times 10^9\text{ M}^{-1}\text{ s}^{-1}$  [63]). Lee


et al. [19] found no formation of DMA during the reaction, suggesting that DMA is not a degradation by-product of NDMA by hydroxyl radicals.

Pathway 2



Dosing H<sub>2</sub>O<sub>2</sub> (~3 mg/L) has shown to have a negligible effect on the degradation of NDMA [27<sup>•</sup>, 49]. A previous study [16] reported that H<sub>2</sub>O<sub>2</sub> addition prior to UV irradiation enhanced NDMA degradation by only 10%, while another pilot-scale study [56] reported a negligible impact of H<sub>2</sub>O<sub>2</sub> addition (0–10 mg/L) on NDMA degradation. Inefficient degradation of NDMA by the powerful HO<sup>•</sup> oxidant may stem from a slow reaction of the methylene carbon-centred radical with dissolved oxygen to form a peroxy radical (Eq. 10) [62–64].

High concentrations of H<sub>2</sub>O<sub>2</sub> can lead to a reduction in the rate of NDMA degradation due to the scavenging of hydroxyl radicals ( $k_{\text{HO}^\bullet} = 2.7 \times 10^7\text{ M}^{-1}\text{ s}^{-1}$ ) [65, 66]:



Hydroxyl radicals can also react with natural organic matter in the source waters, which reduces the availability of hydroxyl radicals for reaction with NDMA [64]. Other inorganic and organic compounds in water can also scavenge hydroxyl radicals and reduce the rate of NDMA degradation. The hydroxyl radical scavengers include carbonate, bicarbonate, nitrite and bromide [65, 67]. Other parameters that influence NDMA degradation include NDMA concentrations, UV intensity, solution pH and anion concentrations [54].

UV lamps with lower wavelengths are preferable due to the larger molar adsorption coefficient of H<sub>2</sub>O<sub>2</sub> at shorter wavelength. Nevertheless, the effects of UV lamp selection on the destruction of NDMA are not conclusive. For example, Sharpless et al. [49] reported a 30% increase in the fluence-based rate constant with the addition of 100 mg/L of H<sub>2</sub>O<sub>2</sub> under LP UV irradiation, while effect from a MP UV lamp was negligible. In contrast, a pilot-scale UV lamp study [56] revealed that the addition of 5 and 10 mg/L H<sub>2</sub>O<sub>2</sub> had a negligible impact, but the results from a MP UV lamp slightly enhanced NDMA degradation.




## Degradation of NDMA by UV/TiO<sub>2</sub> AOP

Photocatalysts typically used for water treatment include TiO<sub>2</sub> [68, 69]. UV irradiation of heterogeneous TiO<sub>2</sub> generates

hydroxyl radicals (HO<sup>•</sup>) and singlet oxygen (<sup>1</sup>O<sub>2</sub>) at the surface through multiple pathways [70]. The photocatalytic reaction is initiated by UV irradiation of TiO<sub>2</sub>. The semiconductor has two energy bands: an energy band occupied by free electrons (valence band (VB)) and an energy band that is generally empty (conduction band (CB)). The VB is lower in energy than the CB. The energy difference between the VB and the CB is called the bandgap. When electrons in the VB absorb UV that has a sufficient photon energy (*hν*) greater than the energy of the bandgap (e.g. λ = <390 nm for anatase TiO<sub>2</sub>), the electrons are excited and jump to the CB forming electron-hole pairs—positive holes in the VB (h<sup>+</sup><sub>TiO<sub>2</sub></sub>) and negative electrons in the CB (e<sup>-</sup><sub>TiO<sub>2</sub></sub>) (Eq. 14). Water molecules that come in contact with the positive holes react to form hydroxyl radicals and protons (H<sup>+</sup>) (Eq. 15), while dissolved oxygen reacts with electrons to form superoxide ions (O<sub>2</sub><sup>-</sup>) (Eq. 16). The dissolved oxygen concentration is a key parameter in the photocatalytic degradation of chemical contaminants in UV/TiO<sub>2</sub> AOP [71].






<sup>1</sup>O<sub>2</sub> is formed through the oxidation of O<sub>2</sub><sup>-</sup> on h<sup>+</sup><sub>TiO<sub>2</sub></sub> in the VB (Eq. 17). <sup>1</sup>O<sub>2</sub> can also be formed through the reaction of O<sub>2</sub><sup>-</sup> with another O<sub>2</sub><sup>-</sup> (Eq. 18), H<sub>2</sub>O<sub>2</sub> (Eqs. 19 and 20) and HO<sup>•</sup> (Eq. 21) [72, 73].





Photocatalytic degradation of NDMA by hydroxyl radicals is initiated with the attack at one of three positions: methyl group, amine nitrogen or nitrosyl nitrogen, as proposed by Lee et al. [74]. Attack of the methyl group (pathway A) leads to the formation of a carbon-centred NDMA radical (<sup>•</sup>CH<sub>2</sub>(CH<sub>3</sub>)N-N=O), which reacts with oxygen molecule and forms peroxy radical (<sup>•</sup>O<sub>2</sub>CH<sub>2</sub>(CH<sub>3</sub>)N-N=O) (Eq. 22). The peroxy radical degrades to alkoxy radical intermediates (Eq. 23) and eventually forms MA (Eq. 24) [75, 76]. This is the major pathway under oxygen-rich conditions.

Pathway A





With the attack of amine nitrogen (pathway B), protonated *N*-methylidenemethylamine (CH<sub>2</sub>=N<sup>+</sup>HCH<sub>3</sub>) is generated, which decays to MA and HCOH by hydrolysis (Eq. 25). A recent study conducted by Guo et al. [77] suggested another pathway for NDMA degradation that is initiated with the attack of the same amine nitrogen but formation of DMA (Eq. 26).

Pathway B



With the attack of nitrosyl nitrogen (pathway C), NDMA is converted into dimethylaminy radical ((CH<sub>3</sub>)<sub>2</sub>N<sup>•</sup>) and nitric acid (Eq. 27) [74]. The dimethylaminy radical, which is unreactive with oxygen but reactive with H-atom-donating compounds, decays to DMA through e<sup>-</sup>/H<sup>+</sup> or an H-atom abstraction from NDMA (Eq. 28) [50, 78].

Pathway C




The resultant MA and DMA can be adsorbed onto TiO<sub>2</sub> where DMA can be degraded into MA, ammonium (NH<sub>4</sub><sup>+</sup>) and NO<sub>3</sub><sup>-</sup>, while MA further degraded into NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> [79].

In addition to the formation of hydroxyl radicals, <sup>1</sup>O<sub>2</sub> generated through UV/TiO<sub>2</sub> AOP [80, 81] can play an important role in the photocatalytic degradation of NDMA. NDMA reacts with <sup>1</sup>O<sub>2</sub> and favourably degrades to DMA through the pathway described in Eq. 29 [25]. The specific location of <sup>1</sup>O<sub>2</sub> attack among three positions (i.e. methyl group, amine nitrogen or nitrosyl nitrogen) has not been clarified.

Pathway D



Pathways of photocatalytic degradation of NDMA are summarised in Fig. 1. The proportion of formed MA and DMA varies depending on the solution matrix (e.g. pH) and the characteristics of photocatalysts [25]. For example, the concentration of formed MA can be two to four times greater than that of formed DMA with pure TiO<sub>2</sub>, while with surface-coated TiO<sub>2</sub> DMA can be formed at concentrations higher than MA [74]. In



**Fig. 1** Pathways of NDMA destruction by UV/TiO<sub>2</sub> AOP

addition to DMA and MA, the photocatalytic destruction of NDMA leads to the formation of NO<sub>2</sub><sup>-</sup> and NO<sub>3</sub><sup>-</sup>, whose concentrations can be as much as formed DMA and MA [77•]. The photocatalytic destruction of NDMA in a high pH solution (e.g. pH = >10) can also form ammonium ions [74].

Recent advances in materials science have allowed the use of TiO<sub>2</sub> with a different morphology for NDMA removal. A notable example is a study conducted by Guo et al. [25••] where TiO<sub>2</sub> was prepared in the form of TiO<sub>2</sub> nanotubes (diameter = 5–6 nm) from anatase TiO<sub>2</sub> to attain surface area larger than commercial TiO<sub>2</sub> powders and particles (e.g. Degussa P25 [48, 51] and Aeroxide P25 [26]). Their study revealed that <sup>1</sup>O<sub>2</sub> formed with the TiO<sub>2</sub> nanotubes contributed to NDMA degradation greater than typical TiO<sub>2</sub> particles. Guo et al. [25••] speculated that the enhanced photocatalytic degradation of NDMA with smaller TiO<sub>2</sub> particles could be attributed to small pores where NDMA molecules could enter. This could also be attributed to greater formation of <sup>1</sup>O<sub>2</sub> due to the increased number of adsorbed oxygen molecules with a larger surface [80].

Photocatalytic degradation of chemicals by UV/TiO<sub>2</sub> AOP could be improved with surface modifications or doping of the TiO<sub>2</sub> heterogeneous photocatalyst [82]. For example, a previous study [74] reported that silica-loaded TiO<sub>2</sub> with UV irradiation can enhance NDMA destruction by 30% as compared to unmodified TiO<sub>2</sub>. Another study by Guo et al. [77•] modified the surface of TiO<sub>2</sub> nanotubes with gold nanoparticles. Interestingly, NDMA destruction with these TiO<sub>2</sub> nanotubes led to the formation of DMA as a major by-product. This indicates that the selection of additives could alter the major pathways of NDMA degradation (pathways A–D) and alter the formation of major by-products. Doping TiO<sub>2</sub> with different materials is an effective strategy to modify the optical properties of TiO<sub>2</sub> that leads to an increase in the absorption of the visible (Vis) light (λ > 380 nm). TiO<sub>2</sub>

particles doped with metals or non-metal dopants attain new energy levels between their VB and CB, which reduce the bandgap and allow Vis light to be absorbed [83]. Many previous studies have demonstrated the effects of Vis light on the photocatalytic degradation of contaminants such as endocrine-disrupting compounds and pharmaceuticals [70, 84]. Nevertheless, no attempts have been reported in regard to NDMA destruction by UV-Vis/TiO<sub>2</sub> AOP.

## Summaries and Future Roadmap

The UV photolysis and UV-based AOPs discussed in this review are summarised in Table 1. This review focused on a fundamental UV technology (i.e. direct UV), a commercialised AOP (i.e. UV/H<sub>2</sub>O<sub>2</sub>) and a potential AOP without continuous chemical addition (i.e. UV/TiO<sub>2</sub>). UV photolysis is very effective for NDMA degradation, but a log reduction of NDMA typically requires a UV dose ≥1000 mJ/cm<sup>2</sup>. In addition, UV alone is ineffective for 1,4-dioxane removal, which requires an AOP with radical generation (e.g. hydroxyl) for effective destruction. Therefore, UV/H<sub>2</sub>O<sub>2</sub> AOP has been employed at most of the recent potable water facilities. However, this AOP requires continuous dosing of H<sub>2</sub>O<sub>2</sub> and often requires quenching to remove the residual. The UV/TiO<sub>2</sub> AOP technology can be a good alternative to direct photolysis and UV/H<sub>2</sub>O<sub>2</sub> AOP as recent studies have shown promising performance for NDMA degradation, and it does not require continuous chemical dosing. The technology relies solely on UV irradiation of the TiO<sub>2</sub> heterogeneous photocatalyst for the formation of hydroxyl radicals and singlet oxygen.

One of the uncertainties of the UV/TiO<sub>2</sub> AOP technology includes the frequency of TiO<sub>2</sub> surface cleaning. Although hydroxyl radicals can essentially destroy most organic compounds, the adsorption of impurities—both organics and inorganics—onto TiO<sub>2</sub> could occur over long-term operations screening the surface of the photocatalyst from UV light. Moreover, the lifetime of the catalyst is of great interest as it contributes to the operational costs. Short catalyst lifetime renders the feasibility of UV/TiO<sub>2</sub> AOP questionable. To scale up the technology, further investigation is necessary for photocatalytic reactor design. Most UV/TiO<sub>2</sub> AOP systems used for laboratory- and pilot-scale studies apply immobilised TiO<sub>2</sub> beds or TiO<sub>2</sub> fine particles in a slurry form [68]. Immobilised TiO<sub>2</sub> beds must be in direct contact with UV irradiation, and much of the surface area remains inaccessible and inactive. A slurry system can utilise large

**Table 1** Comparison of UV (photolysis) and UV-based advanced oxidation technologies for NDMA removal

	Advantages	Disadvantages	Technology level [61]
UV	Well established	Not effective for 1,4-dioxane	Commercial
UV/H <sub>2</sub> O <sub>2</sub>	Well established	Requirement of continuous chemical addition	Commercial
UV/TiO <sub>2</sub>	No continuous chemical addition	Not sufficiently established	Research to commercial



surface active sites, while it has operational difficulties in a separation system to avoid the loss of TiO<sub>2</sub> fine particles for reconditioning/recycling and reuse. Overall, a feasibility study focusing on the continuous use of TiO<sub>2</sub> and photocatalytic reactor design is required to consider UV/TiO<sub>2</sub> AOP as an alternative to the most commonly used AOP in PR—UV/H<sub>2</sub>O<sub>2</sub> AOP.

## Conclusions

This paper comprehensively reviewed UV-based advanced oxidation technologies currently used and available for NDMA removal. Direct photolysis at wavelengths below 270 nm is a very effective way of NDMA removal. However, depending on pH, direct photolysis produces DMA as a major by-product. Since DMA is a major NDMA precursor, regeneration of NDMA can occur if chloramine is used after the UV treatment. UV/H<sub>2</sub>O<sub>2</sub> AOP—the most common approach that is based on UV irradiation in combination with H<sub>2</sub>O<sub>2</sub> dosing—does not provide a significant improvement in NDMA removal in comparison with the direct photolysis. UV/TiO<sub>2</sub> AOP is an alternative approach that does not involve continuous dosing of H<sub>2</sub>O<sub>2</sub> (or other oxidants) and subsequent quenching step. Hydroxyl radicals and singlet oxygen are formed by UV/TiO<sub>2</sub> AOP, both of which are generated at the catalytic surface through UV irradiation. NDMA destruction by UV/TiO<sub>2</sub> AOP forms both DMA and MA, at ratios that vary depending on the reaction conditions and surface properties of TiO<sub>2</sub>. Further improvement of NDMA degradation can be expected by developing the surface morphology modification and doping methods of TiO<sub>2</sub>. In addition, further research on other uncertainties (e.g. the lifetime of TiO<sub>2</sub>, cleaning methods and photocatalytic reactor design) is also needed to scale up the technology in PR applications.

**Acknowledgments** The authors gratefully acknowledge Dr. Kenneth P. Ishida from Orange County Water District (CA, USA) for his assistance in the preparation of this manuscript.

## Compliance with Ethical Standards

**Conflict of Interest** The authors declare that they have no conflict of interest.

## References

Papers of particular interest, published recently, have been highlighted as:

- Of importance,
- Of major importance

1. Mitch WA, Sharp JO, Trussell RR, Valentine RL, Alvarez-Cohen L, Sedlak DL. *N*-nitrosodimethylamine (NDMA) as a drinking water contaminant: a review. *Environ Eng Sci*. 2003;20(5):389–404. doi:10.1089/109287503768335896.
2. Drewes JE, Khan SJ. Contemporary design, operation, and monitoring of potable reuse systems. *J Water Res Desalination*. 2015;5(1):1–7. doi:10.2166/wrd.2014.148.
3. USEPA. *N*-Nitrosodimethylamine (CASRN 62-75-9) (Accessed December 16, 2016)1993.
4. Gerrity D, Pecson B, Trussell RS, Trussell RR. Potable reuse treatment trains throughout the world. *J Water Supply Res*. 2013;62(6):321–38. doi:10.2166/aqua.2013.041.
5. Farré MJ, Döderer K, Hearn L, Poussade Y, Keller J, Gernjak W. Understanding the operational parameters affecting NDMA formation at advanced water treatment plants. *J Hazard Mater*. 2011;185(2–3):1575–81. doi:10.1016/j.jhazmat.2010.10.090.
6. Libotean D, Giral J, Rallo R, Cohen Y, Giral F, Ridgway HF, et al. Organic compounds passage through RO membranes. *J Membr Sci*. 2008;313(1–2):23–43. doi:10.1016/j.memsci.2007.11.052.
7. Fujioka T, Khan SJ, Poussade Y, Drewes JE, Nghiem LD. *N*-nitrosamine removal by reverse osmosis for indirect potable water reuse—a critical review based on observations from laboratory-, pilot- and full-scale studies. *Sep Purif Technol*. 2012;98(0):503–15. doi:10.1016/j.seppur.2012.07.025.
8. Zenker MJ, Borden RC, Barlaz MA. Occurrence and treatment of 1,4-dioxane in aqueous environments. *Environ Eng Sci*. 2003;20(5):423–32. doi:10.1089/109287503768335913.
9. Environment Protection and Heritage Council, National Health and Medical Research Council, Natural Resource Management Ministerial Council. Australian guidelines for water recycling: managing health and environmental risks (phase 2) - augmentation of drinking water supplies. Canberra: Government of Australia; 2008.
10. Linge KL, Blair P, Busetti F, Rodriguez C, Heitz A. Chemicals in reverse osmosis-treated wastewater: occurrence, health risk, and contribution to residual dissolved organic carbon. *J Water Supply Res*. 2012;61(8):494–505. doi:10.2166/aqua.2012.047.
11. California Department of Public Health. Drinking water programs, certificates and licenses, 1, 4-dioxane notification level. <http://www.waterboards.ca.gov>. (accessed 1 December 2016).
12. Fujioka T, Khan SJ, McDonald JA, Roux A, Poussade Y, Drewes JE, et al. *N*-nitrosamine rejection by reverse osmosis membranes: a full-scale study. *Water Res*. 2013;47(16):6141–8. doi:10.1016/j.watres.2013.07.035.
13. Swaim PDM, Rory; Mueller, Paul; Vorissis, Mary; Erdal, Ufuk; Carter, William. Implementing an effective UV advanced oxidation process. *Proceedings of the Water Environment Federation, Disinfection*. 2009:678–88. Doi:10.2175/193864709793847555.
14. Ishida KP, Cooper WJ. Analysis of parameters affecting process efficiency, energy consumption, and carbon footprint in water reuse. Alexandria, VA: WaterReuse Research Foundation; 2015.
15. Keen O, Dotson A, Linden K. Evaluation of hydrogen peroxide chemical quenching agents following an advanced oxidation process. *J Environ Eng*. 2012;139(1):137–40. doi:10.1061/(ASCE)EE.1943-7870.0000619.
16. Kruithof JC, Kamp PC, Martijn BJ. UV/H<sub>2</sub>O<sub>2</sub> treatment: a practical solution for organic contaminant control and primary disinfection. *Ozone-Sci Eng*. 2007;29(4):273–80. doi:10.1080/01919510701459311.
17. Stepien DK, Diehl P, Helm J, Thorns A, Puttmann W. Fate of 1,4-dioxane in the aquatic environment: from sewage to drinking water. *Water Res*. 2014;48:406–19. doi:10.1016/j.watres.2013.09.057.
18. Xu B, Chen Z, Qi F, Ma J, Wu F. Inhibiting the regeneration of *N*-nitrosodimethylamine in drinking water by UV photolysis combined with ozonation. *J Hazard Mater*. 2009;168(1):108–14. doi:10.1016/j.jhazmat.2009.02.014.
19. Lee C, Yoon J, Von Gunten U. Oxidative degradation of *N*-nitrosodimethylamine by conventional ozonation and the advanced

- oxidation process ozone/hydrogen peroxide. *Water Res.* 2007;41(3):581–90. doi:10.1016/j.watres.2006.10.033.
20. Pisarenko AN, Stanford BD, Yan D, Gerrity D, Snyder SA. Effects of ozone and ozone/peroxide on trace organic contaminants and NDMA in drinking water and water reuse applications. *Water Res.* 2012;46(2):316–26. doi:10.1016/j.watres.2011.10.021.
  21. Deng Y, Ezyske CM. Sulfate radical-advanced oxidation process (SR-AOP) for simultaneous removal of refractory organic contaminants and ammonia in landfill leachate. *Water Res.* 2011;45(18):6189–94. doi:10.1016/j.watres.2011.09.015.
  22. Wang Y, Le Roux J, Zhang T, Croué J-P. Formation of brominated disinfection byproducts from natural organic matter isolates and model compounds in a sulfate radical-based oxidation process. *Environ Sci Technol.* 2014;48(24):14534–42. doi:10.1021/es503255j.
  23. Watts MJ, Hofmann R, Rosenfeldt EJ. Low-pressure UV/Cl<sub>2</sub> for advanced oxidation of taste and odor. *Journal of the American Water Works Association (JAWWA).* 2012;104(1):58–65. doi:10.5942/jawwa.2012.104.0006.
  24. Deng Y, Zhao R. Advanced oxidation processes (AOPs) in wastewater treatment. *Current Pollution Reports.* 2015;1(3):167–76. doi:10.1007/s40726-015-0015-z. **This article provides an overview of advanced oxidation processes used in wastewater treatment that facilitates understanding of the oxidation theory.**
  25. Guo X, Li Q, Zhang M, Long M, Kong L, Zhou Q, et al. Enhanced photocatalytic performance of *N*-nitrosodimethylamine on TiO<sub>2</sub> nanotube based on the role of singlet oxygen. *Chemosphere.* 2015;120:521–6. doi:10.1016/j.chemosphere.2014.09.002. **This article assessed photocatalytic degradation performance of N-nitrosodimethylamine by TiO<sub>2</sub> nanotube. It also identified the role of singlet oxygen as well as hydroxyl radicals during the photocatalytic degradation.**
  26. Alvarez-Corena JR, Bergendahl JA, Hart FL. Advanced oxidation of five contaminants in water by UV/TiO<sub>2</sub>: reaction kinetics and byproducts identification. *J Environ Manag.* 2016;181:544–51. doi:10.1016/j.jenvman.2016.07.015.
  27. Sgroi M, Roccaro P, Oelker GL, Snyder SA. *N*-nitrosodimethylamine (NDMA) formation at an indirect potable reuse facility. *Water Res.* 2015;70:174–83. doi:10.1016/j.watres.2014.11.051. **This article focused on NDMA precursors in RO permeate. They identified that NDMA formation during chloramination of the RO permeate.**
  28. Shen R, Andrews SA. Demonstration of 20 pharmaceuticals and personal care products (PPCPs) as nitrosamine precursors during chloramine disinfection. *Water Res.* 2011;45(2):944–52. doi:10.1016/j.watres.2010.09.036.
  29. Le Roux J, Gallard H, Croué J-P. Chloramination of nitrogenous contaminants (pharmaceuticals and pesticides): NDMA and halogenated DBPs formation. *Water Res.* 2011;45(10):3164–74. doi:10.1016/j.watres.2011.03.035.
  30. Mitch WA, Schreiber IM. Degradation of tertiary alkylamines during chlorination/chloramination: implications for formation of aldehydes, nitriles, halonitroalkanes, and nitrosamines. *Environ Sci Technol.* 2008;42(13):4811–7. doi:10.1021/es703017z.
  31. Mitch WA, Gerecke AC, Sedlak DL. A *N*-nitrosodimethylamine (NDMA) precursor analysis for chlorination of water and wastewater. *Water Res.* 2003;37(15):3733–41. doi:10.1016/S0043-1354(03)00289-6.
  32. Pehlivanoglu-Mantas E, Hawley EL, Deeb RA, Sedlak DL. Formation of nitrosodimethylamine (NDMA) during chlorine disinfection of wastewater effluents prior to use in irrigation systems. *Water Res.* 2006;40(2):341–7. doi:10.1016/j.watres.2005.11.012.
  33. Schreiber IM, Mitch WA. Influence of the order of reagent addition on NDMA formation during chloramination. *Environ Sci Technol.* 2005;39(10):3811–8. doi:10.1021/es0483286.
  34. Soltermann F, Lee M, Canonica S, von Gunten U. Enhanced *N*-nitrosamine formation in pool water by UV irradiation of chlorinated secondary amines in the presence of monochloramine. *Water Res.* 2013;47(1):79–90. doi:10.1016/j.watres.2012.09.034.
  35. Poussade Y, Roux A, Walker T, Zavlanos V. Advanced oxidation for indirect potable reuse: a practical application in Australia. *Water Sci Technol.* 2009;60(9):2419–24. doi:10.2166/wst.2009.665.
  36. Drewes JE, Khan SJ. Chapter 16: Water reuse for drinking water augmentation. In: Edzwald JK, editor. *Water quality & treatment: a handbook on drinking water.* 6th ed. New York: McGraw-Hill; 2011. p. 16.1–16.48.
  37. Plumlee M, López Mesas M, Heidlberger A, Ishida K, Reinhard M. *N*-nitrosodimethylamine (NDMA) removal by reverse osmosis and UV treatment and analysis via LC–MS/MS. *Water Res.* 2008;42(1–2):347–55. doi:10.1016/j.watres.2007.07.022.
  38. Yangali-Quintanilla V, Maeng SK, Fujioka T, Kennedy M, Amy G. Proposing nanofiltration as acceptable barrier for organic contaminants in water reuse. *J Membr Sci.* 2010;362(1–2):334–45. doi:10.1016/j.memsci.2010.06.058.
  39. USEPA. 1,4-Dioxane (CASRN 123–91-1) (Accessed December 16, 2016)1993.
  40. Nawrocki J, Andrzejewski P. Nitrosamines and water. *J Hazard Mater.* 2011;189(1–2):1–18. doi:10.1016/j.jhazmat.2011.02.005.
  41. Sedlak D, Kavanaugh M. Removal and destruction of NDMA and NDMA precursors during wastewater treatment. *Water Reuse Foundation;* 2006.
  42. Krauss M, Longrée P, Dorusch F, Ort C, Hollender J. Occurrence and removal of *N*-nitrosamines in wastewater treatment plants. *Water Res.* 2009;43(17):4381–91. doi:10.1016/j.watres.2009.06.048.
  43. Krasner SW, Mitch WA, McCurry DL, Hanigan D, Westerhoff P. Formation, precursors, control, and occurrence of nitrosamines in drinking water: a review. *Water Res.* 2013;47(13):4433–50. doi:10.1016/j.watres.2013.04.050.
  44. Yoon S, Nakada N, Tanaka H. Occurrence and removal of NDMA and NDMA formation potential in wastewater treatment plants. *J Hazard Mater.* 2011;190(1–3):897–902. doi:10.1016/j.jhazmat.2011.04.010.
  45. Andrzejewski P, Kasprzyk-Hordern B, Nawrocki J. *N*-nitrosodimethylamine (NDMA) formation during ozonation of dimethylamine-containing waters. *Water Res.* 2008;42(4–5):863–70. doi:10.1016/j.watres.2007.08.032.
  46. Schreiber IM, Mitch WA. Nitrosamine formation pathway revisited: the importance of chloramine speciation and dissolved oxygen. *Environ Sci Technol.* 2006;40(19):6007–14. doi:10.1021/es060978h.
  47. Farré MJ, Keller J, Holling N, Poussade Y, Gernjak W. Occurrence of *N*-nitrosodimethylamine precursors in wastewater treatment plant effluent and their fate during ultrafiltration-reverse osmosis membrane treatment. *Water Sci Technol.* 2011;63(4):605–12. doi:10.2166/wst.2011.207.
  48. Lee C, Choi W, Yoon J. UV photolytic mechanism of *N*-nitrosodimethylamine in water: roles of dissolved oxygen and solution pH. *Environ Sci Technol.* 2005;39(24):9702–9. doi:10.1021/es051235j.
  49. Sharpless C, Linden K. Experimental and model comparisons of low- and medium-pressure Hg lamps for the direct and H<sub>2</sub>O<sub>2</sub> assisted UV photodegradation of *N*-nitrosodimethylamine in simulated drinking water. *Environ Sci Technol.* 2003;37(9):1933–40. doi:10.1021/es025814p.
  50. Stefan M, Bolton J. UV direct photolysis of *N*-nitrosodimethylamine (NDMA): kinetic and product study. *Helv Chim Acta.* 2002;85(5):1416–26. doi:10.1002/1522-2675(200205)85:5<1416::AID-HLCA1416>3.0.CO;2-I.
  51. Lee C, Choi W, Kim Y, Yoon J. UV photolytic mechanism of *N*-nitrosodimethylamine in water: dual pathways to methylamine versus dimethylamine. *Environ Sci Technol.* 2005;39(7):2101–6. doi:10.1021/es0488941.
  52. Chow YL, Lau MP, Perry RA, Tam JNS. Photoreactions of nitroso compounds in solution. XX. Photoreduction, photoelimination, and

- photoaddition of nitrosamines. *Can J Chem.* 1972;50(7):1044–50. doi:10.1139/v72-164.
53. Plumlee MH, Reinhard M. Photochemical attenuation of *N*-nitrosodimethylamine (NDMA) and other nitrosamines in surface water. *Environ Sci Technol.* 2007;41(17):6170–6. doi:10.1021/es070818l.
  54. Zhou C, Gao N, Deng Y, Chu W, Rong W, Zhou S. Factors affecting ultraviolet irradiation/hydrogen peroxide (UV/H<sub>2</sub>O<sub>2</sub>) degradation of mixed *N*-nitrosamines in water. *J Hazard Mater.* 2012;231–232:43–8. doi:10.1016/j.jhazmat.2012.06.032.
  55. Xu B, Chen Z, Qi F, Shen J, Wu F. Factors influencing the photodegradation of *N*-nitrosodimethylamine in drinking water. *Front Environ Sci Eng China.* 2009;3(1):91–7. doi:10.1007/s11783-009-0013-8.
  56. Lekkerkerker-Teunissen K, Knol AH, Derks JG, Heringa MB, Houtman CJ, Hofman-Caris CHM, et al. Pilot plant results with three different types of UV lamps for advanced oxidation. *Ozone-Sci Eng.* 2013;35(1):38–48. doi:10.1080/01919512.2013.721317.
  57. Sakai H, Kosaka K, Takizawa S. Degradation of *N*-nitrosodimethylamine by mercury-free excimer UV lamps. *Environ Eng Sci.* 2016;33(5):341–6. doi:10.1089/ees.2015.0328.
  58. Sakai H, Takamatsu T, Kosaka K, Kamiko N, Takizawa S. Effects of wavelength and water quality on photodegradation of *N*-nitrosodimethylamine (NDMA). *Chemosphere.* 2012;89(6):702–7. doi:10.1016/j.chemosphere.2012.06.025.
  59. Zhang J-Y, Boyd IW. Lifetime investigation of excimer UV sources. *Appl Surf Sci.* 2000;168(1–4):296–9. doi:10.1016/S0169-4332(00)00628-0.
  60. Goldstein S, Aschengrau D, Diamant Y, Rabani J. Photolysis of aqueous H<sub>2</sub>O<sub>2</sub>: quantum yield and applications for polychromatic UV actinometry in photoreactors. *Environ Sci Technol.* 2007;41(21):7486–90. doi:10.1021/es071379t.
  61. Collins J, Bolton JR. Advanced oxidation handbook. American Water Works Association; 2016.
  62. Mezyk S, Cooper W, Madden K, Bartels D. Free radical destruction of *N*-nitrosodimethylamine in water. *Environ Sci Technol.* 2004;38(11):3161–7. doi:10.1021/es0347742.
  63. Mezyk SP, Ewing DB, Kiddle JJ, Madden KP. Kinetics and mechanisms of the reactions of hydroxyl radicals and hydrated electrons with nitrosamines and nitramines in water. *J Phys Chem A.* 2006;110(14):4732–7. doi:10.1021/jp056017r.
  64. Landsman NA, Swancutt KL, Bradford CN, Cox CR, Kiddle JJ, Mezyk SP. Free radical chemistry of advanced oxidation process removal of nitrosamines in water. *Environ Sci Technol.* 2007;41(16):5818–23. doi:10.1021/es070275f.
  65. Wols BA, Hofman-Caris CHM. Review of photochemical reaction constants of organic micropollutants required for UV advanced oxidation processes in water. *Water Res.* 2012;46(9):2815–27. doi:10.1016/j.watres.2012.03.036.
  66. Muruganandham M, Swaminathan M. Photochemical oxidation of reactive azo dye with UV–H<sub>2</sub>O<sub>2</sub> process. *Dyes Pigments.* 2004;62(3):269–75. doi:10.1016/j.dyepig.2003.12.006.
  67. Yang Y, Pignatello JJ, Ma J, Mitch WA. Effect of matrix components on UV/H<sub>2</sub>O<sub>2</sub> and UV/S<sub>2</sub>O<sub>8</sub><sup>2-</sup> advanced oxidation processes for trace organic degradation in reverse osmosis brines from municipal wastewater reuse facilities. *Water Res.* 2016;89:192–200. doi:10.1016/j.watres.2015.11.049.
  68. Chong MN, Jin B, Chow CWK, Saint C. Recent developments in photocatalytic water treatment technology: a review. *Water Res.* 2010;44(10):2997–3027. doi:10.1016/j.watres.2010.02.039.
  69. Park H, Choi W. Effects of TiO<sub>2</sub> surface fluorination on photocatalytic reactions and photoelectrochemical behaviors. *J Phys Chem B.* 2004;108(13):4086–93. doi:10.1021/jp036735i.
  70. Lazar M, Varghese S, Nair S. Photocatalytic water treatment by titanium dioxide: recent updates. *Catalysts.* 2012;2(4):572. doi:10.3390/catal2040572.
  71. Zhang D, Qiu R, Song L, Eric B, Mo Y, Huang X. Role of oxygen active species in the photocatalytic degradation of phenol using polymer sensitized TiO<sub>2</sub> under visible light irradiation. *J Hazard Mater.* 2009;163(2–3):843–7. doi:10.1016/j.jhazmat.2008.07.036.
  72. Daimon T, Nosaka Y. Formation and behavior of singlet molecular oxygen in TiO<sub>2</sub> photocatalysis studied by detection of near-infrared phosphorescence. *J Phys Chem C.* 2007;111(11):4420–4. doi:10.1021/jp070028y.
  73. Atlante A, Passarella S. Detection of reactive oxygen species in primary cultures of cerebellar granule cells. *Brain Res Protocol.* 1999;4(3):266–70. doi:10.1016/S1385-299X(99)00028-8.
  74. Lee J, Choi W, Yoon J. Photocatalytic degradation of *N*-nitrosodimethylamine: a mechanism, product distribution, and TiO<sub>2</sub> surface modification. *Environ Sci Technol.* 2005;39(17):6800–7. doi:10.1021/es0481777.
  75. Maillard B, Ingold KU, Scaiano JC. Rate constants for the reactions of free radicals with oxygen in solution. *J Am Chem Soc.* 1983;105(15):5095–9. doi:10.1021/ja00353a039.
  76. Alfassi ZB, Mosseri S, Neta P. Halogenated alkylperoxyl radicals as oxidants: effects of solvents and of substituents on rates of electron transfer. *J Phys Chem.* 1987;91(12):3383–5. doi:10.1021/j100296a056.
  77. Guo X, Shao H, Kong L, Long M, Zhang M, Zhou Q, et al. Probing the effect of nanotubes on *N*-nitrosodimethylamine photocatalytic degradation efficiency and reaction pathway. *Chem Eng Sci.* 2016;144:1–6. doi:10.1016/j.ces.2016.01.018. **This article evaluated the effects of modification of TiO<sub>2</sub> nanotube on the photocatalytic degradation of *N*-nitrosodimethylamine and identified dimethylamine as a major by-product.**
  78. Wink D, Desrosiers M. Unusual spin-trap chemistry for the reaction of hydroxyl radical with the carcinogen *N*-nitrosodimethylamine. *International Journal of Radiation Applications and Instrumentation Part C.* 1991;38(5):467–72. doi:10.1016/1359-0197(91)90063-8.
  79. Helali S, Puzenat E, Perol N, Safi M-J, Guillard C. Methylamine and dimethylamine photocatalytic degradation—adsorption isotherms and kinetics. *Appl Catal A Gen.* 2011;402(1–2):201–7. doi:10.1016/j.apcata.2011.06.004.
  80. Daimon T, Hirakawa T, Kitazawa M, Suetake J, Nosaka Y. Formation of singlet molecular oxygen associated with the formation of superoxide radicals in aqueous suspensions of TiO<sub>2</sub> photocatalysts. *Appl Catal A Gen.* 2008;340(2):169–75. doi:10.1016/j.apcata.2008.02.012.
  81. Hirakawa T, Yawata K, Nosaka Y. Photocatalytic reactivity for O<sub>2</sub><sup>-</sup> and OH radical formation in anatase and rutile TiO<sub>2</sub> suspension as the effect of H<sub>2</sub>O<sub>2</sub> addition. *Appl Catal A Gen.* 2007;325(1):105–11. doi:10.1016/j.apcata.2007.03.015.
  82. Henderson MA. A surface science perspective on photocatalysis. *Surf Sci Rep.* 2011;66(6–7):185–297. doi:10.1016/j.surfrep.2011.01.001.
  83. Pelaez M, Nolan NT, Pillai SC, Seery MK, Falaras P, Kontos AG, et al. A review on the visible light active titanium dioxide photocatalysts for environmental applications. *Appl Catal B Environ.* 2012;125:331–49. doi:10.1016/j.apcatb.2012.05.036.
  84. Fagan R, McCormack DE, Dionysiou DD, Pillai SC. A review of solar and visible light active TiO<sub>2</sub> photocatalysis for treating bacteria, cyanotoxins and contaminants of emerging concern. *Mater Sci Semicond Process.* 2016;42(Part 1):2–14. doi:10.1016/j.mssp.2015.07.052.